

CP Violation and FCNC in the Third Family from Effective Supersymmetry

Ann E. Nelson

*Department of Physics 1560, University of Washington, Seattle, WA 98195-1560,
USA*

ABSTRACT

I discuss a “more minimal” modification of the minimal supersymmetric standard model, in which supersymmetry breaking is connected with the physics of flavor. Flavor Changing Neutral Currents (FCNC) for the first two families are suppressed, and for the third family may be of comparable size to the FCNC in the Standard Model.

Modifications of the Minimal Standard Model are haunted by a fundamental dichotomy¹: namely that in most extensions, Flavor Changing Neutral Current (FCNC) constraints are naturally satisfied only if the new physics scale is above 10–1000 TeV, whereas natural electroweak symmetry breaking requires new physics below ~ 1 TeV. In table 1 I contrast the virtues and omissions of the Standard Model with two popular extensions: the Minimal Supersymmetric Standard Model (MSSM) with soft supersymmetry breaking, and Technicolor. On the left I have listed various experimental observations that a good theory should explain. It is clear that all these models have many shortcomings, however each of them explains at least one experimental fact in a way that is so beautiful it is hard to believe nature would not make use of it. Now I would like to discuss what features of each of these models makes them so successful, and how one might put them together in a single theory.²

1. The observed spectrum and couplings

Both the Standard Model and the MSSM are very succesful at accomodating the observed fermion masses, mixings, and couplings to the weak gauge bosons, and the key to this success is a fundamental Higgs scalar.

2. Naturally small M_W/M_{pl}

Here we should borrow solutions from both the MSSM and from technicolor. Weak scale supersymmetry naturally stabilizes the value of the weak scale against perturbative quantum corrections,³ and Dynamical Supersymmetry Breaking⁴ (DSB) can explain why this scale is so far below the Planck scale, in a manner analogous to the dynamical electroweak symmetry breaking of Technicolor.⁵

Table 1

	Standard Model	Minimal Supersymmetric Standard Model	Technicolor
<i>Spectrum, Precision EW measurements</i>			
M_W/M_{pl} } <i>Value</i>			
} <i>Perturbative stability</i>			
<i>B conservation</i>			
<i>L conservation</i>			
<i>FCNC</i>			
<i>Weak CP</i>			
<i>Strong CP</i>			
<i>Cosmological constant</i>			
<i>Quantum gravity</i>			

3. Baryon Conservation, Lepton Conservation, FCNC, and weak CP violation

Here the most compelling solution is found in the Standard Model—gauge invariance automatically requires all dangerous operators to be of dimension higher than 4 and so irrelevant. Baryon and Lepton number conservation arise as “accidental” symmetries, and FCNC are suppressed by approximate global flavor symmetries which are also accidental. Unfortunately this is not at all true in the MSSM, where global symmetries must be imposed by hand.

4. The Strong CP Problem, the Cosmological Constant Problem, and Quantum Gravity

None of these issues are addressed by any models for physics at the weak scale, and it is likely that only a theory which is valid for experimentally inaccessible distance scales will explain them.

5. Effective Supersymmetry

Our conclusion from the above critique is that we should include both elementary scalars and supersymmetry in our theory of electroweak symmetry breaking. In addition, *a new gauge interaction* is desirable for two reasons:

1. to dynamically break supersymmetry
2. to remove the possibility of low dimension B and L violating operators by ensuring that any dangerous gauge invariant operators are *automatically* sufficiently irrelevant.

It is most economical if the same gauge interaction does both jobs, so the DSB gauge interaction also couples directly to at least some ordinary fields. Such a gauge interaction can give a large (hopefully positive) supersymmetry breaking mass to the scalar components of the superfields it couples to. If the scalar masses are very large, ~ 20 TeV, for the first two generations of squarks and sleptons, the FCNC problems of supersymmetry are alleviated via a new mechanism, *squark and slepton decoupling*.^{2,6}

The top squark and left handed bottom squark couple strongly to the Higgs and natural electroweak symmetry breaking requires that these be lighter than 1 TeV. Therefore they should not carry any new supersymmetry breaking gauge interactions. Having these sparticles be light does not cause any (so far) problematic FCNC. By treating the generations differently, the dichotomy between the high scale necessary for FCNC suppression and the lower scale determined by natural electroweak symmetry breaking is removed.

Note that all dangerous dimension 4 and 5 baryon violating operators of the MSSM involve at least one superfield from the first two generations, so the proton stability could be guaranteed by new gauge symmetries for the first two generations alone.

Automatic suppression of dimension 3 and 4 lepton number violating interactions requires that the down-type Higgs and the sleptons be distinguished by different gauge interactions as well. One possibility is for the down type Higgs to carry the new supersymmetry breaking gauge interaction, which would make the scalar very heavy and lead to naturally large $\tan \beta$. Another possibility is for all generations of left handed leptons to carry the new interaction and so all the left handed sleptons would be heavy.

Such a new interaction could also suppress the couplings of the first two generations to the Higgs, explaining the generational mass hierarchy. Thus, at least in principle, a new gauge interaction carried by the first two generations of quarks and leptons could solve a remarkable number of problems. It could produce the gauge hierarchy via DSB, while suppressing FCNC, SUSY contributions to particle electric dipole moments, and baryon and lepton number violation. It could even explain the fermion mass hierarchy. Unfortunately no completely explicit and successful examples exist, but several recently constructed models come close.^{2,7}

Of course we have to assume that new gauge symmetries for ordinary particles are spontaneously broken or confined above some scale Λ . The scale of heavy

<i>observed phenomenon</i>	<i>Standard Model</i>	<i>SUSY (universality)</i>	<i>SUSY(alignment)</i>	<i>Effective SUSY</i>
See $D\bar{D}$ mixing	X	X	predicts	possible
No $D\bar{D}$ mixing	predicts	predicts	X	possible
V_{td} , Δm_B not SM rel.	X	possible	possible ($<30\%$)	possible
δ_{KM} , ε not SM rel.	X	possible	possible	possible
Non-SM CP phase relations in B_d mixing	X	X	$<\pi/10$	possible, $O(\pi/2)$
Non-SM CP phase relations in B_s mixing	X	X	X	possible, $< O(\pi/2)$
Non-SM CP phase relations in direct B decays	X	X	X	possible, $O(\pi)$

Table 2

squark and slepton masses will then be of order $\Lambda_S^2/\Lambda \sim 20$ TeV, where Λ_S is the fundamental supersymmetry breaking scale. The ordinary gauginos, top squarks and left handed bottom squark, which only couple weakly to the supersymmetry breaking sector, are significantly lighter, below 1 TeV. Other scalars of the third generation may be either light or heavy.

The presence of a light left handed bottom squark will enhance the SUSY box diagram contribution to $B - \bar{B}$ mixing. In addition there is no reason to expect the SUSY phase in this contribution to be the same as that of the standard model diagram. Similarly, there can be significant SUSY penguins, with new CP violating phases, for the third generation. Thus the B factory signals for Effective Supersymmetry can be quite different from those in the MSSM with universal squark masses, or with squark masses aligned with the down quark masses.⁸ In Table 2 I summarize the potentially observable differences between the different SUSY models. Clearly the framework I have described is not sufficiently predictive to be ruled out by experiments at low energies. However I am optimistic that hints of physics beyond the Standard Model will be found in such experiments. A decisive test will have to wait for the next generation of collider experiments.

Discovery of a squark or slepton from the first two generations rules out Effective Supersymmetry, however we expect the Higgs, top squarks, left handed bottom squark, Higgsinos and gauginos to be found.

Acknowledgements

I would like to thank the organizers for inviting me to this most interesting conference, and David Kaplan for the use of the tables done on his NEXT. This work was supported in part by the DOE under grant no. #DE-FG03-96ER40956.

References

1. L. J. Hall, V. A. Kostelecky, S. Raby, Nucl. Phys. B267, 415 (1986); H. Georgi, Phys. Lett. 169B, 231 (1986);
2. A. G. Cohen, D. B. Kaplan, A. E. Nelson, Phys. Lett. B388, 588 (1996)
3. S. Dimopoulos, H. Georgi, Nucl. Phys. B193, 150 (1981); N. Sakai, Z. Phys. C11, 153 (1981)
4. E. Witten, Nucl. Phys. B188, 513 (1981)
5. S. Weinberg, Phys. Rev. D13, 974 (1976), *ibid.* D19, 1277 (1979); L. Susskind, Phys. Rev. D20, 2619 (1979)
6. M. Dine, A. Kagan, S. Samuel, Phys. Lett. B243, 250 (1990); S. Dimopoulos, G. F. Giudice, Phys. Lett. B357, 573 (1995); A. Pomarol, D. Tommasini, Nucl. Phys. B466 3 (1996)
7. G. Dvali, A. Pomarol, CERN-TH-96-192, hep-ph/9607383; P. Binetruy, E. Dudas, Phys. Lett. B389, 503 (1996); R.N. Mohapatra, A. Riotto, Phys. Rev. D55, 1138 (1997); A. E. Nelson, D. Wright UW-PT-97-03, hep-ph/9702359
8. A. G. Cohen, D. B. Kaplan, F. Lepeintre, A. E. Nelson, Phys. Rev. Lett. 78, 2300 (1997)